



- Conigliera

General information

Basic information

Wetland location:	Marine/Coastal
Wetland type:	Wetland system
Natural / Artificial:	Natural
Hydrological interaction with other wetland:	No -
Water salinity:	Salty (> 18.0 g/l)
Fresh water entry:	Catchment area (precipitation)
Open water area (%):	> 95
Hydroperiod:	Temporary/Intermittent

Geographic information

Governorate:	Monastir
Island:	Petite Kuriat, Conigliera
Coordinates (WGS84):	11.006907 E - 35.768649 N

Wetland condition

Wetland condition:	1 - Untouched. No signs of man-made changes
---------------------------	---

Ramsar wetland types

Ramsar type	Coverage (%)
A -- Permanent shallow marine waters in most cases less than six metres deep at low tide; includes sea bays and straits	5 - 25
E -- Sand, shingle or pebble shores; includes sand bars, spits and sandy islets; includes dune systems and humid dune slacks	5 - 25
H -- Intertidal marshes; includes salt marshes, salt meadows, saltings, raised salt marshes; includes tidal brackish and freshwater marshes	5 - 25
J -- Coastal brackish/saline lagoons; brackish to saline lagoons with at least one relatively narrow connection to the sea	51 - 75

Property status

Public

Protection statuses & other designations

Ecosystem Services, Activities & Impacts

Ecosystem Services

Type of Ecosystem service	Ecosystem service	Scale of Benefit	Importance
Cultural services	Recreation and tourism		
Provisioning services	Food		
Regulatory services	Flood hazard regulation		
Regulatory services	Storm hazard regulation		
Regulatory services	Water purification		
Regulatory services	Water regulation		
Supporting services	Provision of habitat		

Activities on wetland

Activities	Intensity
020 = Resource conservation	High
030 = Species conservation	High
210 = Professional fishing	High
240 = Taking / Removal of fauna general	High
690 = Other leisure and tourism impacts not referred to above	High

Activities on drainage basin

Activities	Intensity
020 = Resource conservation	High
030 = Species conservation	High
210 = Professional fishing	High
240 = Taking / Removal of fauna general	High
690 = Other leisure and tourism impacts not referred to above	High

Impacts

Impact type	Intensity
WSR = Salt water intrusion of surface water	High

Habitats & Vegetation

Species

Flora

Species	Dominance	Reference
Arthrocnemum macrostachyum (Moric.) K. Koch		
Cakile maritima Scop.		
Echium arenarium Guss.		
Euphorbia paralias L.		
Euphorbia terracina L.		
Frankenia laevis L.		
Halocnemum strobilaceum (Pall.) M. Bieb.		
Hordeum marinum Huds.		
Lamarckia aurea (L.) Moench		
Limoniastrum monopetalum (L.) Boiss.		
Rostraria cristata (L.) Tzvelev		
Sarcocornia fruticosa (L.) A.J. Scott		

Sarcocornia perennis (Mill.) A.J. Scott subsp. Per

Suaeda vera Forssk. ex J.F. Gmel.

Suaeda vermiculata Forssk. ex J.F. Gmel.

Fauna

Fishes	Presence in wetland	References
<i>Carcharodon carcharias</i> (Linnaeus, 1758)		
<i>Cethorinus maximus</i> (Gunnerus, 1765)		
<i>Isurus oxyrinchus</i> (Rafinesque, 1810)		
<i>Lamna nasus</i> (Bonnaterre, 1788)		

References

Representative Image & Map